

PULSAR PSR B0656+14, THE MONOGEM RING, AND THE ORIGIN OF THE ‘KNEE’ IN THE PRIMARY COSMIC RAY SPECTRUM

S. E. THORSETT,¹ R. A. BENJAMIN,² WALTER F. BRISKEN,³ A. GOLDEN,⁴ & W. M. GOSS³

To appear in the Astrophys. J. Lett.

ABSTRACT

The Monogem ring is a bright, diffuse, 25°-diameter supernova remnant easily visible in soft X-ray images of the sky. Projected within the ring is a young radio pulsar, PSR B0656+14. An association between the remnant and pulsar has been considered, but was seemingly ruled out by the direction and magnitude of the pulsar proper motion and by a distance estimate that placed the pulsar twice as far from Earth as the remnant. Here we show that in fact the pulsar was born very close to the center of the expanding remnant, both in distance and projection. The inferred pulsar and remnant ages are in good agreement. The conclusion that the pulsar and remnant were born in the same supernova explosion is nearly inescapable. The remnant distance and age are in remarkable concordance with the predictions of a model for the primary cosmic ray energy spectrum in which the ‘knee’ feature is produced by a single dominant source.

Subject headings: pulsars: individual (PSR B0656+14)—supernova remnants—acceleration of particles

1. INTRODUCTION

A bright soft X-ray enhancement in the Monoceros and Gemini constellations was first detected in rocket experiments over thirty years ago (Bunner et al. 1971), and resolved into a shell-like structure by *HEAO-2* a decade later (Nousek et al. 1981). Spectral studies with *ROSAT* confirmed that this so-called ‘Monogem ring’ is a supernova remnant, probably in the adiabatic expansion phase (Plucinsky et al. 1996). The distance is poorly constrained by evolutionary arguments—self-consistent Sedov-Taylor models were found by Plucinsky et al. (1996) at all distances between 100 and 1300 pc—but distances around 300 pc are preferred on grounds of supernova energetics. At this distance, the model age is 86,000 years, explosion energy is 1.9×10^{50} erg, and current radius is 66 pc. The inferred interstellar medium density is $5 \times 10^{-3} \text{ cm}^{-3}$, typical of the hot interstellar medium.

The radio pulsar PSR B0656+14 lies very close in projection to the center of the Monogem ring, and an association between the objects seems natural (Córdova et al. 1989; Nousek et al. 1981; Thompson et al. 1991). The pulsar is young, with a characteristic spin-down timescale of 110 kyr (Taylor et al. 1993), pulsed non-thermal optical and X-ray emission, and unpulsed thermal X-ray emission consistent with a 10^5 year-old cooling neutron star Finley et al. (1993). But because the pulsar distance was estimated from interstellar dispersion measurements (Taylor et al. 1993) to be 760 pc—more than twice the best estimate for the remnant—and a proper motion measurement appeared to show the pulsar moving towards the center of the remnant (Thompson & Córdova




FIG. 1.— The Monogem ring, as seen in the *ROSAT* all-sky survey in the 0.25–0.75 keV x-ray band. PSR B0656+14 is marked with cross-hairs, and the 9.2° circle centered on this point shows the primary ring structure. The estimated position of the pulsar 10^5 yrs ago is marked with a small square. J2000 coordinates are shown, and the Galactic plane is indicated by a diagonal line. The ring is imperfect: there is an apparent blow-out to the west, at high galactic latitude, and a missing quadrant to the northwest, perhaps due to foreground absorption or to slower expansion into a dense region. The bright source at 06:17 +22:34 is the SNR IC443.

1994), a physical association has been widely regarded as unlikely (Kaspi 1998, for example). New very long baseline interferometric measurements (Briskin et al. 2003) lead us to reconsider this negative conclusion. The distance, now accurately known from parallax to be 288^{+33}_{-27} pc, is much lower than previously believed, and the proper motion, 44 mas/yr, is about 40% smaller. As we will show, these measurements, together with a re-examination of the X-ray data, convincingly demonstrate that the pulsar and remnant were born in a single supernova explosion about 10^5 yrs ago.

2. THE PULSAR-REMNANT ASSOCIATION

2.1. Agreement in projection

¹ Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064;

² Department of Physics, University of Wisconsin, Madison, WI 53706;

³ National Radio Astronomy Observatory, PO Box O, Socorro, NM 87801;

⁴ National University of Ireland, Newcastle Road, Galway, Republic of Ireland;

The Monogem ring (Fig. 1) shows significant deviations from circular symmetry, with what may be a blow-out to the west, away from the Galactic plane, and a missing quadrant in the northwest, where foreground absorption is relatively high (Plucinsky et al. 1996). Because of the lack of symmetry, the morphological center of the ring is somewhat uncertain. For example, at low energy (e.g., the *ROSAT* R1 band image displayed as Fig. 6 in Plucinsky et al. 1996), the portion of the ring in the galactic plane is not readily visible, making the remnant appear smaller and offset slightly away from the plane; the arc is evident in higher energy observations (Fig. 1, and also note Fig. 8 of Plucinsky et al. 1996), suggesting the importance of greater absorption on lines of sight near the plane.

In projection, the pulsar, at $(l, b) \sim (201.1^\circ, +8.3^\circ)$, lies well within the ring. To test the consistency of its position with the morphological center of the ring, we have fitted ‘by eye’ a circle to the incomplete ring, centered on the pulsar. This fit, shown in Fig. 1, demonstrates that the pulsar’s position is not inconsistent with the morphological center of the ring, within admittedly large uncertainties.

With the measured proper motion, 44 mas/yr, the pulsar has moved about 1.25° in 10^5 yrs, from roughly $(l, b) \sim (200.5^\circ, +7.2^\circ)$. This inferred birth position appears slightly further from the current center of the ring, but not unacceptably so. It is also important to distinguish the morphological center of the remnant from its true dynamical center: we should expect an old remnant expanding into a medium with density decreasing away from the galactic plane to have its apparent center offset from its explosion center in the direction away from the plane, just as observed (Hnatyk & Petruk 1999).

In the end, we conclude that the pulsar position at birth was within a few degrees of the current morphological center of the ring. We note that the *a priori* likelihood of an unrelated background source projected within the ring falling this close to the center is about 5%. However, on its own the positional coincidence is a strong argument neither for nor against an association.

2.2. Agreement in distance

The strongest argument against a physical association between the pulsar and remnant has been the pulsar distance, estimated at 760 pc from the measured column density of free electrons. Sedov modeling formally allows a wide range of distances for the Monogem ring, but a distance of about 300 pc has been preferred. Although estimates of pulsar distances through interstellar dispersion are fairly crude (the quoted uncertainty was 25% (Taylor et al. 1993)) the distance discrepancy appeared significant. The small parallax distance, 288^{+33}_{-27} pc, thus came as a surprise. (We note, though, that the greater distance implied an unacceptably large thermal X-ray luminosity: an attempt to directly estimate the distance from X-ray modeling yielded ~ 200 pc or less (Golden & Shearer 1999).)

With this precise pulsar distance available, we consider whether the distance estimate for the remnant can also be improved. A strong interstellar O VI absorption feature at +105 km/s has been detected in the star 15 Mon, in the southern region of the ring (Jenkins et al.

1998). The parallax distance of 15 Mon is 313^{+93}_{-58} pc (Perryman et al. 1997). Because O VI at such high velocity is rarely found in the disk, and the velocity is similar to that expected from a remnant at this stage, we consider this to be a secure upper limit to the distance of the Monogem ring. Other evidence suggests the ring is no farther than the pulsar. The absorption is low, $N_H \lesssim 1.0 \times 10^{20} \text{ cm}^{-2}$ (Plucinsky et al. 1996), compared to $\sim 1.5 \times 10^{20} \text{ cm}^{-2}$ for the pulsar (Edelstein et al. 2000). The pulsar dispersion measure is much higher than expected at a distance of 290 pc, suggesting a line of sight through a highly ionized region.

It is also unlikely that the remnant is significantly closer than the pulsar. At an assumed distance of 300 pc, the inferred explosion energy is already low: 1.9×10^{50} erg. A distance below 230 pc would imply an implausibly low energy, below 10^{50} erg. We conclude that the pulsar and remnant distances agree to within ~ 50 pc. Indeed, since the shell radius is ~ 70 pc, the evidence is strong that the pulsar is currently within the expanding supernova shell.

2.3. Agreement in age

Finally, we must consider the relative ages of the pulsar and remnant. The remnant age, from Sedov modeling, has been estimated at 86,000 yrs for a distance of 300 pc. Two lines of evidence suggest a comparable age for the neutron star. First, its current temperature is in good agreement with standard cooling models for a 10^5 yr old neutron star (van Riper et al. 1995). Second, the characteristic spin-down timescale for the pulsar is $P/2\dot{P} = 1.1 \times 10^5$ yrs.

The second argument deserves more comment. The age of a pulsar slowing with a torque proportional to a constant power n of its frequency, $\dot{f} \propto -f^n$, can be expressed as

$$\tau = \frac{P}{(n-1)\dot{P}} \left[1 - \left(\frac{P_0}{P} \right)^{n-1} \right], \quad (1)$$

where P_0 is the initial spin period. For magnetic-dipole braking, the ‘braking index’ $n = 3$, so if the pulsar is born spinning much faster than its current period, $\tau \approx P/2\dot{P}$, while the age is overestimated if the pulsar were born near its current period.

For a handful of very young pulsars, where the braking index can be measured directly, $n \lesssim 3$ (Briskin et al. 2003). For middle-aged pulsars, the good agreement between timing ages and so-called kinetic ages ($z/\dot{z}$, where z is the height above the Galactic plane) requires a mean braking index near 3 (Bailes 1989; Bhattacharya et al. 1992; Lorimer et al. 1997). We acknowledge that the timing age can be misleading. Several putative pulsar-SNR associations suggest large discrepancies between timing and true ages: for example, PSR J1012–5226, in the ~ 7 kyr old remnant PKS 1209–51/52, has a spin-down age of $\sim 200 - 900$ kyr (Pavlov et al. 2002), and PSR J1811–1925, with a spin-down age of 24 kyr, has been associated with the SNR of AD 386 (Torii et al. 1999). Nevertheless, a recent comparison of kinetic and timing ages for 21 pulsars under 10 Myr in age found a typical discrepancy of only $\sim 40\%$ (Briskin et al. 2003). We conclude that both the timing and cooling ages of the pulsar are consistent with the remnant age.

2.4. Summary of arguments for an association

The angular position, distance, and age of the pulsar and remnant are all in excellent agreement with the hypothesis that both were born at the same location, ~ 300 pc away, 10^5 yrs ago. The likelihood that this is merely a positional and temporal coincidence is extremely low. Crudely estimating the Galactic Type II supernova rate at one per century and the region within 300 pc of Earth as $(300 \text{ pc}/10 \text{ kpc})^2 \approx 0.1\%$ of the area of the Galactic disk, the mean supernova interval in this region is about 100 kyr. The chance that two unrelated supernovae occur within ~ 25 kyr, within 5° on the sky, and within 50 pc in radial distance is vanishingly small. Perhaps the most persuasive argument for an association comes from assuming the opposite. If this 10^5 yr old remnant is not that of the supernova that formed the 10^5 yr old pulsar, then where is the remnant of that supernova? Many old remnants are of course invisible because of distance or environment, but if the Monogem ring and pulsar were formed in different supernovae then they occurred in very close physical and temporal proximity. The Monogem ring itself is evidence that a remnant of age 10^5 years expanding into this particular environment is visible at 300 pc. We conclude that a single supernova, 300 pc away and a hundred thousand years ago, formed both PSR B0656+14 and the Monogem ring.

3. THE MONOGEN RING AS A ‘SINGLE SOURCE’ FOR PEV COSMIC RAYS

Supernova remnants are generally believed to be acceleration sites for cosmic rays (Berezhko 1996; Berezhko & Ksenofontov 1999; Blandford & Ostriker 1980; Shklovsky 1953), and in this case it is intriguing to note that the Monogem ring may hold a clue to a long-standing puzzle: the origin of the steepening in the primary cosmic-ray spectrum at about 3×10^{15} eV (3 PeV), called the ‘knee.’

Between about 10^{10} and a few times 10^{18} eV, the differential cosmic ray energy spectrum is well-described by a broken power law, proportional to below the knee and above the knee. (A recent review can be found in Wefel (2003).) Possible explanations for the knee are extremely varied, including loss of the most energetic particles from the Galaxy, unknown physical processes in the development of the atmospheric shower through which the cosmic rays are detected, or, most likely, a termination in the acceleration process. A typical cut-off energy for ac-

celeration in supernova remnants is $Z \times 10^{14}$ eV, where Z is the nuclear charge, though the exact cut-off will vary by perhaps an order of magnitude with variations in the explosion energy, magnetic field, interstellar medium density, and age (Berezhko 1996; Erlykin & Wolfendale 1997; Legage & Cesarsky 1983).

Recently, Erlykin and Wolfendale (EW) have drawn attention (Erlykin & Wolfendale 1997) to the sharpness of the knee feature, which has been a challenge for models in which the break arises from propagation effects or from a stochastic superposition of multiple sources with varying high-energy cut-offs. EW propose that around the knee a single nearby source dominates the cosmic ray spectrum, which is otherwise a smooth superposition of the contributions from many supernova remnants throughout the Galaxy and from whatever sources provide the higher energy cosmic rays, up to 10^{20} eV and beyond. At energies of a few PeV, this single source alone produces 60% of the flux at Earth. (This possibility has also recently been considered by other authors (Berezhko & Ksenofontov 1999, for example).) Although the data are not yet conclusive, EW have identified the knee as most likely due to oxygen nuclei, with a smaller second break at about 10 PeV due to iron nuclei.

The shape and amplitude of the knee feature have led EW to predict (Erlykin & Wolfendale 2003) that the single source is a 90–100 kyr old supernova remnant between 300 and 350 pc from Earth, expanding in an under-dense medium. The match with the properties of the Monogem ring is striking, though almost certainly in part coincidental considering the large remaining uncertainties in cosmic ray acceleration and diffusion models. Nevertheless, an important objection to the single source model for the knee feature has been the lack of a suitable source (Bhadra 2002). That objection now appears to have been removed.

S.E.T. is supported by the NSF under grant AST-0098343, R.A.B. is supported by NASA ATP grant NAG5-12128, and A.G. is supported by Enterprise Ireland under grant SC/2001/322. This research has made use of the *ROSAT* all-sky survey data, which have been processed at MPE, and of the telescopes of the National Radio Astronomy Observatory, a facility of the NSF operated under cooperative agreement by Associated Universities, Inc. We thank an anonymous referee for suggestions that improved the manuscript.

REFERENCES

- Bailes, M. 1989, *ApJ*, 342, 917
 Berezhko, E. G. 1996, *Astropart. Phys.*, 5, 367
 Berezhko, E. G. & Ksenofontov, L. T. 1999, *JETP*, 89, 391
 Bhadra, A. 2002, *J. Phys. G: Nucl. Part. Phys.*, 28, 397
 Bhattacharya, D., Wijers, R. A. M. J., Hartman, J. W., & Verbunt, F. 1992, *A&A*, 254, 198
 Blandford, R. D. & Ostriker, J. P. 1980, *ApJ*, 237, 793
 Brisen, W. F., Fruchter, A. S., Goss, W. M., Herrnstein, R. S., & Thorsett, S. E. 2003, *Astron. J.*, submitted
 Bunner, A. N., Coleman, P. L., Kraushaar, W. L., & McCammon, D. 1971, *ApJ*, 167, L3
 Córdoba, F. A., Hjellming, R. M., Mason, K. O., & Middleditch, J. 1989, *ApJ*, 345, 451
 Edelstein, J., Seon, K. I., Golden, A., & Min, K. W. 2000, *ApJ*, 539, 902
 Erlykin, A. D. & Wolfendale, A. W. 1997, *J. Phys. G: Nucl. Part. Phys.*, 23, 979
 —. 2003, *J. Phys. G: Nucl. Part. Phys.*, 29, 709
 Finley, J. P., Ögelman, H., & Kisiloglu, U. 1993, *ApJ*, 394, L21
 Golden, A. & Shearer, A. 1999, *A&A*, 342, L5
 Hnatyk, B. & Petruk, O. 1999, *A&A*, 344, 295
 Jenkins, E. B., Sofia, U. S., & Sonneborn, G. 1998, in *The Hot Universe*, ed. K. Koyama, S. Kitamoto, & M. Itoh (Dordrecht: Kluwer Academic), 273–274
 Kaspi, V. M. 1998, *Adv. Space Res.*, 21, 167
 Legage, P. O. & Cesarsky, C. J. 1983, *A&A*, 125, 249
 Lorimer, D. R., Bailes, M., & Harrison, P. A. 1997, *MNRAS*, 289, 592
 Nousek, J. A., Cowie, L. L., Hu, E., Linblad, C. J., & Garmire, G. P. 1981, *ApJ*, 248, 152

- Pavlov, G. G., Zavlin, V. E., Sanwal, D., & Trümper, J. 2002, *ApJ*, 569, L95
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., Hoeg, E., Bastian, U., Bernacca, P. L., Crézé, M., Donati, F., Grenon, M., van Leeuwen, F., van der Marel, H., Mignard, F., Murray, C. A., Le Poole, R. S., Schrijver, H., Turon, C., Arenou, F., Froeschlé, M., & Petersen, C. S. 1997, *A&A*, 323, L49
- Plucinsky, P. P., Snowden, S. L., Aschenbach, B., Egger, R., Edgar, R. J., & McCammon, D. 1996, *ApJ*, 463, 224
- Shklovsky, I. S. 1953, *Dokl. Akad. Nauk. USSR*, 91, 475
- Taylor, J. H., Manchester, R. N., & Lyne, A. G. 1993, *ApJS*, 88, 529
- Thompson, R. J. & Córdova, F. A. 1994, *ApJ*, 421, L13
- Thompson, R. J., Córdova, F. A., Hjellming, R. M., & Fomalont, E. B. 1991, *ApJ*, 366, L83
- Torii, K., Tsunemi, H., Dotani, T., Mitsuda, K., Kawai, N., Kinugasa, K., Saito, Y., & Shibata, S. 1999, *ApJ*, 523, L69
- van Riper, K. A., Link, B., & Epstein, R. I. 1995, *ApJ*, 448, 294
- Wefel, J. P. 2003, *J. Phys. G: Nucl. Part. Phys.*, 29, 821